# **\* NARCAN** (naloxone HCI) **NASAL SPRAY**

Attachment B

# **QUICK START GUIDE Opioid Overdose Response Instructions**

### Use NARCAN Nasal Spray (naloxone hydrochloride) for known or suspected opioid overdose in adults and children.

Important: For use in the nose only.

Do not remove or test the NARCAN Nasal Spray until ready to use.

Identify Opioid Overdose and Check for Response

**Ask** person if he or she is okay and shout name.

Shake shoulders and firmly rub the middle of their chest.

#### Check for signs of opioid overdose:

• Will not wake up or respond to your voice or touch

**Remove** NARCAN Nasal Spray from the box.

- Breathing is very slow, irregular, or has stopped
- Center part of their eye is very small, sometimes called "pinpoint pupils"
- Lay the person on their back to receive a dose of NARCAN Nasal Spray.

Peel back the tab with the circle to open the NARCAN Nasal Spray.



Call for

medical

Evaluate,

Support

help,

and

emergency

**Hold** the NARCAN nasal spray with your thumb on the bottom of the plunger and your first and middle fingers on either side of the nozzle.

## Gently insert the tip of the nozzle into either nostril.

• Tilt the person's head back and provide support under the neck with your hand. Gently insert the tip of the nozzle into one nostril, until your fingers on either side of the nozzle are against the bottom of the person's nose.

Press the plunger firmly to give the dose of NARCAN Nasal Spray. • Remove the NARCAN Nasal Spray from the nostril after giving the dose.

Get emergency medical help right away.

Move the person on their side (recovery position) after giving NARCAN Nasal Spray.

Watch the person closely.

If the person does not respond by waking up, to voice or touch, or breathing normally another dose may be given. NARCAN Nasal Spray may be dosed every 2 to 3 minutes, if available.

# Repeat Step 2 using a new NARCAN Nasal Spray to

© 2015 ADAPT Pharma, Inc. NARCAN® is a registered trademark licensed to ADAPT Pharma Operations Limited, A1009

give another dose in the other nostril. If additional NARCAN Nasal Sprays are available, repeat step 2 every 2 to 3 minutes until the person responds or emergency medical help is received.















